



**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

ESU

The  
**Patent  
Office**

PCT/GB 99/04326  
09/868394

20 DECEMBER 1999

INVESTOR IN PEOPLE

GB99/4326

EPO - DG 1

20. 01. 2000

(59)

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ


REC'D 16 FEB 2000	
WIPO	PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *Andrew Gurney*

Dated 20 December 1999




21DEC98 E413355-1 002029  
P01/7700 0.00 - 9928087.8

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office  
Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference



2. Patent application number

(The Patent Office will fill in his part)

**9828087.8**

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

SMITHKLINE BEECHAM PLC  
NEW HORIZONS COURT, BRENTFORD,  
MIDDLESEX TW8 9EP

UNITED KINGDOM

*5200 974002*

4. Title of the invention

Novel Method

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent  
(including the postcode)

Patents ADP number (if you know it)

CORPORATE INTELLECTUAL PROPERTY

SMITHKLINE BEECHAM PLC  
TWO NEW HORIZONS COURT  
BRENTFORD  
MIDDLESEX TW8 9EP

*5200 974004*

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country      Priority application number      Date of filing  
(if you know it)      (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application      Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer yes if:

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is named as an applicant, or
  - c) any named applicant is a corporate body
- See note (d)

9. Enter the number of sheets for any of the following items you are filing with this form.  
Do not count copies of the same document

Continuation sheets of this form	0
Description	7
Claim(s)	0
Abstract	0
Drawings	0

10. If you are also filing any of the following, state how many against each item.

**Priority Documents**

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 1/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(please specify)

11.

We request the grant of a patent on the basis of this application

Signature K Rutter Date 18-Dec-98

12. Name and daytime telephone number of person to contact in the United Kingdom

K Rutter 01279 644396

**Warning**

*After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- For details of the fee and ways to pay please contact the Patent Office.*

## Novel Method

5 The invention relates to a novel method and more particularly to a method for the detection of compounds that mimic, potentiate or inhibit the physiological effects of the ob-protein

The ob-protein (or leptin) is a secreted hormone that acts as signal from adipose tissue to other organs to regulate weight and energy balance (Zhang et. al., *Nature*, 1994, **372**, 425). Additional roles for the ob-protein in haematopoietic and reproductive function have been suggested (Cioffi et. al. *Nature Medicine*, 1996, **2**(5), 585). Protein molecules that contain a core composed of four  $\alpha$ -helices forming a bundle of up-up-down-down topology comprise a family of cytokines and growth factors. Proteins of this family cause homo- and hetero-oligomerisation of membrane receptors known to activate kinase cascades resulting in gene transcription. Receptors of the family which are activated by oligomerisation fall into two broad classes; those such as epidermal growth factor receptor, which possess integral tyrosine kinase activity in their intracellular domains (A. Ullrich & J. Schlessinger, *Cell*, 1990, **61**, 203-212), and those such as the IL4 and erythropoietin receptors, which lack this activity and mediate their response by way of an associated protein tyrosine kinase (J.N. Ihle et al., *TIBS*, 1994, **19**, 222-227). Both receptor subtypes are activated by cytokines, but the 4-helix bundle proteins activate only the non-integral tyrosine kinase subtype. The non-integral protein tyrosine kinase receptors generally act through a pathway involving Janus kinase (JAK) and their associated signal transducers and activators of transcription (STAT) proteins. On activation STAT proteins bind to DNA response elements thereby controlling gene transcription. Oligonucleotide sequences comprising DNA regulatory elements of the general sequence TT(N)nAA have been identified (Seidel et al., *Proc. Nat. Acad. Sci. USA.*, 1995, **92**, 3041) as STAT response elements. These elements bind STAT proteins in response to signalling molecules such as cytokines.

In copending United Kingdom patent application number 9509164.1 we have described our discovery that the ob-protein is characterised by a four helix bundle tertiary structure. We now believe that the ob-protein interacts with a membrane bound receptor that activates a JAK-STAT kinase cascade and hence forms the basis for an assay system for the detection of compounds that mimic, potentiate or inhibit the physiological effects of the ob-protein. Such an assay has utility in selecting compounds for the treatment of weight, energy balance, haematopoietic, fertility and other disorders modulated by the ob-protein. The assay is especially useful for selecting compounds for the treatment of those disorders related to obesity, anorexia, cachexia and diabetes.

Copending International patent application number PCT/EP96/02291 relates to a novel detection method which uses JAK-STAT technology. We have now found a particularly advantageous detection method which also utilises this technology.

Accordingly, the invention provides a method for the detection of a compound that mimics, potentiates or inhibits the physiological effect of the ob-protein, which method comprises:

(a) for a compound which mimics the physiological effect of the ob-protein, assessing the effect of the compound upon an ob-protein activated signal transducer and activator of transcription (STAT) DNA response element coupled to a reporter gene; or

- 5 (b) for a compound which potentiates or inhibits the physiological effect of the ob-protein, assessing the effect of the compound upon the response provided by ob protein upon an ob-protein activated STAT DNA response element coupled to a reporter gene;

wherein,

- 10 the response element and the reporter are expressed in an ob-protein responsive cell line, which cell line is an endothelium derived cell line.

A suitable source of endothelium-derived cells is a human immortalised endothelial cell line, a murine or other non-human immortalised endothelial cell line, human primary endothelial cells, or murine or other non-human primary endothelial

- 15 cells.

A suitable human endothelium derived cell line is an ECV304-human umbilical cord cell line (see *In Vitro Cell Dev. Biol.* 1990; 26, 265; *In Vitro Cell Dev Biol.* 1991; 27A, 766 or *In Vitro Cell Dev Biol.* 1992; 28A, 380)

A suitable murine endothelial cell line is selected from the list consisting of:

- 20 SVEC4-10 –endothelial lymph node cells, SV40 transformed;  
SVEC4-10EE2 –endothelial lymph node cells, SV40 transformed;  
SVEC-10EHR1 - endothelial lymph node cells, SV40 transformed;  
IP-1B - endothelial lymph node cells, SV40 transformed;  
2F-2B - endothelial lymph node cells, SV40 transformed;  
25 3B-11 - endothelial lymph node cells, SV40 transformed;  
2H-11 –endothelial lymph node cells, SV40 transformed; and  
MS1 (Mile SVEN 1)–endothelial pancreatic islet cells, SV40 transformed.

- For SVE C4-10 see *J. Immunol.* 1990; 144, 521-525; *Am. J. Pathol.* 1991; 139, 743-749; *J. Invest. Dermatol.* 1993; 100, 742-745. SVEC4-10 is the parental cell
- 30 line for a series of endothelial cell lines including; SVEC4-10EE2; SVEC-10HER; 2H-11; 3B-11; 2F-2B; and IP-1B

A suitable human primary endothelial line is selected from the list consisting of:

- 35 HUVEC - human umbilical vein endothelial cells;  
HUAEC - human umbilical artery endothelial cells;  
HAEC – human aortic endothelial cells;  
HPAEC – human pulmonary artery endothelial cells;  
HDMECa – human microvascular endothelial cells, adult dermis; and  
40 HDMECn – human microvascular endothelial cells, neonatal dermis.

All of the above mentioned primary cell lines are commercially available.

A suitable polypeptide which is capable of mediating the stimulation by ob-protein of an ob-protein activated STAT DNA response element is a functional

isoform of the ob-gene receptor, for example that identified in Tartaglia et al., *Cell*, 1995, **83**, 1263.

Suitably, the response element is coupled to a promoter gene, preferably a minimal promoter.

5 A suitable response element is a nucleotide of formula  $TT(N)_n AA$ , where N is any nucleotide and n is 4, 5 or 6.

A favoured response element is selectively activated by the intracellular events mediated the by the ob-protein interacting with its receptor. Such selective response elements can be determined by examining the relative activation of a range of  
10 response element-reporter gene constructs when transfected into an ob-responsive cell line by the ob-protein versus other cytokines.

A favoured response element is a nucleotide of formula  $TT(N)_n AA$ , where N is any nucleotide and n is 4 or 5.

A further suitable response element is TTCCCGGAA.

15 A further suitable response element is that region of the promoter of a gene regulated by the ob-protein that is required for STAT interactions. This gene will depend on the particular therapeutic use of the compounds to be selected by the assay.

A suitable reporter gene is firefly luciferase or chloramphenicol acetyltransferase enzyme.

20 A suitable promoter is a minimal promoter such as the herpes simplex virus thymidine kinase or SV40 promoter.

Other responsive cell lines can be identified using a displacement binding assay. Although binding may not be to a functional long form of the receptor, which is the form that transmits a signal to the cytoplasm. Identification of a functional long  
25 form of the receptor may be by PCR or Northern blot analysis (eg. Human ob-receptor: Tartaglia et al., *Cell*, 1995, **83**, 1263). Ultimately responsive cells are detected by monitoring cellular events in the presence of varying concentrations of leptin. Potential methods for identifying candidate cell lines or monitoring these cellular events include the following:-

30 1. Microphysiometer: This method detects small changes in pH resulting from biochemical changes in the cell. Ob-protein responsive cells upon stimulation may undergo biochemical changes that cause a small change in the extracellular acidification rate which can be detected by a silicon microphysiometer. The microphysiometer biosensor methodology has been reviewed by McConnell, *Science*,  
35 1992, **257**, 1906.

2. Electrophoretic mobility shift assay (EMSA): Nuclear extracts from cells after treatment with ob-protein are mixed with radiolabeled oligonucleotides containing a promiscuous or specific STAT response element DNA sequence. Extracts from cells that respond to the ob-protein may cause a gel shift of the  
40 oligonucleotide for the STAT response element.

References: Book "Recombinant DNA", 2nd Edition, Watson et al. , 1992, Page 158; Lamb et al., *Blood*, 1994, **83**, 2063;

3. Measurement of protein phosphorylation assay: The coupling of receptor activation to the final response through tyrosine phosphorylation of intracellular

proteins may be assayed by the use of antibodies recognising phosphorylated tyrosines. More specifically since the leptin receptor may stimulate tyrosine phosphorylation of the JAK/STAT pathway this method provides a method of detecting leptin response cell lines. Specific JAK/ STAT antibodies may be used alongside antibodies for tyrosine phosphorylation to detect leptin activation in a leptin responsive cell line. Inhibition as well as stimulation of protein phosphorylation may occur. In particular, inhibition by the ob-protein of insulin stimulated phosphorylation of the insulin receptor and insulin receptor substrate-1 has been shown in rat-1 fibroblasts over expressing insulin receptors (Kroder et. al 1996, *Exp. Clin. Endocrinol. Diabetes*, **104**, suppl 2, p66)

4. Displacement binding: After incubation of cell lines with radiolabelled leptin, for example [<sup>125</sup>I]-leptin, the non-specific binding versus specific binding of leptin can be studied by the addition of unlabelled leptin. A high specific to non-specific ratio binding suggests that the cell line may contain the leptin receptor.

5. Detection of the protein for a functional form, preferably a functional long form, of the ob-receptor by use of selective antibodies.

6. Detection of mRNA for a functional form, preferably a functional long form, of the ob-receptor by Northern, RT-PCR or "slot blot" analysis.

7. Detection of increased c-fos mRNA after treatment with leptin. C-fos mRNA may be detected by Northern, RT-PCR or "slot blot" analysis.

Cell lines known to be involved in controlling aspects of the particular disease state for which compounds are being sought are preferred.

Cells lines derived from liver, brain, or pancreatic tissue and fibroblasts are particularly useful for "ob-responsive" cells for the assaying of compounds directed at obesity and diabetes. Certain areas of the brain are the focus of weight controlling and energy balance regulating effects of the ob-protein. The liver controls many metabolic processes that modulate lipid and glucose levels. Cells derived from particular regions of these organs containing the appropriate endogenous JAKs, STAT proteins and other intracellular proteins which are required for mediating the effects of the leptin are preferred.

The response element, the reporter, and preferably the promoter, are suitably incorporated into a vector capable of transfecting the ob-responsive cell line.

Suitable vectors are commercially available vectors, such as pGL2-basic luciferase vector (Promega).

A suitable configuration of the vector is the STAT DNA response element upstream of a promoter and a reporter gene. A more suitable configuration of the vector is the STAT DNA response element in multiple tandem repeats (2-10) upstream of a thymidine kinase promoter and a luciferase reporter gene

Vectors are constructed containing a reporter gene for example firefly luciferase or chloramphenicol acetyltransferase enzyme linked to a minimal promoter for example the herpes simplex virus thymidine kinase or SV40 promoter. The DNA fragments for the STAT response element are inserted into the vector using appropriate restriction enzyme sites upstream of the minimal promoter.



The response element, the reporter and the promoter, as required, are incorporated into the vector using conventional expression techniques, for example the DNA fragments for the response element may be inserted into the vector using appropriate restriction enzyme sites upstream of the minimal promoter.

5       STAT response element-luciferase enzyme reporter systems can be constructed as described by Lamb et al., Blood, 1994, 8, 2063 and Seidel et al., *Proc. Nat. Acad. Sci. USA.*, 1995, 92, 3041.

Ob-responsive cells are transfected with the STAT response element-minimal promoter-luciferase reporter constructs using standard methodology for example the calcium phosphate method (Graham and Van Der Eb, *Virology*, 1973, 52, 456). To correct for differences in transfection efficiency, the cells can be co-transfected with a reference plasmid expressing  $\beta$ -galactosidase activity. After a period of transfection (12-24 hours) the cells are treated with varying concentrations of compound and then harvested and lysed. The lysates are assayed for luciferase, and if appropriate  $\beta$ -galactosidase, activity. Potentiation or antagonist activity can be assayed by pre- or co-addition of an appropriate concentration of ob-protein to the compound under evaluation and measuring the potentiation or reduction in luciferase response relative to that of ob-protein alone. Standard methods exist for assaying luciferase enzyme activity for example Ow et al., *Science*, 1986, 234, 856 and de Wet et al., *Mol. Cell Biol.*, 1987, 7, 725. as well as several commercial kits.

20       Stable cell lines can be generated by transfecting an "ob-responsive" cell line with the reporter construct and a selectable marker. Selectable markers are routinely used to generate stable cell lines as described in Recombinant DNA, 2nd edition, J.D. Watson et. al., 1992, page 216. These stably transfected cell lines can be used to generate high throughput assays for compounds that mimic, potentiate or block the physiological effects of the ob-protein.

The invention also extends to a compound that mimics, potentiates or inhibits the physiological effect of the ob-protein, when identified by the method disclosed herein.

30       The invention also extends to a kit of parts adapted for use in the method disclosed herein.

When used herein 'a compound which mimics the physiological effects of the ob-protein' refers to a compound which is capable of acting in the absence of the ob-protein to either stimulate the ob-protein receptor to provide substantially the same physiological effect as the ob protein or to activate a response down stream of this receptor (post-receptor).

When used herein 'a compound that potentiates the physiological effect of the ob-protein' refers to a compound which enhances the potency and/or maximal physiological effect of the ob-protein.

40       When used herein 'a compound that inhibits the physiological effect of the ob-protein' refers to a compound which reduces or substantially blocks the physiological effect of the ob protein.

The cDNA encoding the functional form of the polypeptide can be transfected under the control of a constitutive promoter, (eg a viral promotor) or a regulatable

5 promoter to optimise the expression of the polypeptide for the identification of agonists or antagonists as necessary. Alternatively, the response element and the reporter are expressed in a cell line, wherein a constitutive or regulatable promoter has been engineered into a position upstream of the chromosomally encoded gene for the ob-protein receptor by the method of homologous recombination. Such methods are reviewed by Waldman, Critical Reviews in Oncology/Hematology, 1992, 12, 49 and a particular example is given in te Riele et al, Proceedings of the National Academy of Sciences, 1992, 89, 5128.

The following examples illustrate the invention but do not limit it in any way.

### Example

#### General Procedure:

- 5 Ob-responsive cells are transfected with a reporter plasmid containing a STAT response element, in multiple tandem copies upstream of a minimal promoter for example herpes simplex thymidine kinase and a luciferase gene reporter construct using standard methodology for example the calcium phosphate method (Graham and Van Der Eb, Virology, 1973, 52, 456). To correct for differences in transfection efficiency, the cells can be co-transfected with a reference plasmid expressing  $\beta$ -galactosidase activity. After a period of transfection (12-24 hours) the cells are treated with varying concentrations of compound and then harvested and lysed. The lysates are assayed for luciferase, and if appropriate  $\beta$ -galactosidase, activity.
- 10 Antagonist activity can be assayed by pre- or co-addition of an appropriate concentration of ob-protein to the compound under evaluation and measuring the reduction in luciferase response relative to that of ob-protein alone. Standard methods exist for assaying luciferase enzyme activity for example Ow et al., *Science*, 1986, 234, 856 and de Wet et al., 1987, 7, 725. as well as several commercial kits.
- 15

#### 20 Example 1

- The particular endothelial cells are transfected with a reporter plasmid, pGL2-basic luciferase vector (Promega) containing an insert of an oligonucleotide corresponding to a four fold tandem repeat of the STAT response element, TTCCCGGAA, upstream of the minimal promoter for herpes simplex thymidine kinase (-35 to +10) using standard methodology for example the calcium phosphate method (Graham and Van Der Eb, Virology, 1973, 52, 456). To correct for differences in transfection efficiency, the cells can be co-transfected with a reference plasmid expressing  $\beta$ -galactosidase activity. After a period of transfection (12-24 hours) the cells are treated with varying concentrations of compound and then harvested and lysed. The lysates are assayed for luciferase, and if appropriate  $\beta$ -galactosidase, activity.
- 25 Antagonist activity can be assayed by pre- or co-addition of an appropriate concentration of ob-protein to the compound under evaluation and measuring the reduction in luciferase response relative to that of ob-protein alone. Standard methods exist for assaying luciferase enzyme activity for example Ow et al., *Science*, 1986, 234, 856 and de Wet et al., *Mol. Cell Biol.*, 1987, 7, 725. as well as several commercial kits.
- 30
- 35

THE PATENT OFFICE  
01 FEB 2000  
NEWPORT